

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/705	A1	(11) International Publication Number: WO 98/53832 (43) International Publication Date: 3 December 1998 (03.12.98)
<p>(21) International Application Number: PCT/US98/10890</p> <p>(22) International Filing Date: 28 May 1998 (28.05.98)</p> <p>(30) Priority Data: 08/866,712 30 May 1997 (30.05.97) US</p> <p>(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US).</p> <p>(72) Inventor: HAUPERT, Garner, T., Jr.; 512 Great Road, Littleton, MA 01460 (US).</p> <p>(74) Agents: ELMORE, Carolyn, S. et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02173 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: USE OF HYPOTHALMIC INHIBITORY FACTOR IN THE TREATMENT OF ISCHEMIC CARDIAC MALFUNCTION</p> <p>(57) Abstract</p> <p>The present invention is based on the unexpected discoveries that hypothalamic inhibitory factor (HIF) has a positive inotropic effect in a whole organ preparation, such as an isolated perfused heart, and that HIF, to a greater extent than ouabain, dilates the coronary arteries in the isolated perfused heart as manifested in increased coronary flow in hearts treated with HIF. The present invention relates to a method for increasing coronary perfusion in a mammalian host comprising administering to the host in need thereof an effective amount of hypothalamic inhibitory factor (HIF). The invention also relates to a method for producing an increased coronary vasodilatory effect in a mammalian host comprising administering to a mammalian host in need thereof an effective amount of HIF. The method of the present invention can be used to prevent and/or treat ischemic cardiac malfunction and coronary artery restenosis.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF HYPOTHALMIC INHIBITORY FACTOR IN THE TREATMENT OF ISCHEMIC CARDIAC MALFUNCTION

BACKGROUND

While heart attack and stroke now cause fewer deaths than previously, deaths due to heart failure (inability of the heart to deliver adequate blood supply to vital organs via the systemic circulation) have increased markedly (82.5%) between 1979 and 1992. In 1997, heart failure is expected to cause at least 250,000 deaths in the United States. More than 400,000 new cases of heart failure are expected to be diagnosed in 1997. In all, it is estimated that more than 4 million Americans are presently living with this dangerous condition.

Sixty to seventy percent of these cases of heart failure are due to ischemic causes, i.e., insufficient delivery of blood flow and vital oxygen via the coronary circulation to the cardiac muscle. This can be due to mechanical obstruction of the coronary arteries, for example by atherosclerotic deposits, to spasm (temporary occlusion) of the coronary arteries, or to restenosis of coronary arteries that have been previously opened by angioplasty (balloon catheter dilation).

Certain segments of the general population are particularly prone to heart failure. Heart attack survivors have a four to six times greater risk of having an additional heart attack. Patients with angina (ischemic coronary pain), diabetes mellitus, or uncontrolled high blood pressure have a two-fold risk of developing heart failure compared to the general population.

Current treatments of heart failure include agents which dilate, and thus, relax blood vessels reducing

-2-

arterial resistance against which the heart must work. Converting enzyme inhibitors such as captopril, enalapril and lisinopril are examples of such arterial vasodilators. Side effects such as rash, persistent cough, excessive
5 lowering of blood pressure and adverse effects on kidney function limit the use of these agents in therapy. Other drugs which work by directly dilating blood vessels (arteries), such as hydralazine, prazosin and doxazosin also reduce the resistance against which the failing heart
10 must pump, but these drugs too can cause excessive lowering of blood pressure, as well as reflex tachycardia (increased heart rate) which increases cardiac work and oxygen demand by the heart muscle already deprived of an adequate oxygen supply in the case of ischemic heart failure.

15 Because of their positive inotropic effect, cardiac glycosides (e.g., digitalis, ouabain) have been considered unrivaled in value for the treatment of heart failure. Positive inotropic effect generally refers to the enhancement of the contractility of the cardiac cells in a
20 dose-dependent manner. Cardiac glycosides are most frequently used therapeutically to increase the adequacy of the circulation in patients with congestive heart failure and to slow the ventricular rate in the presence of atrial fibrillation.

25 However, cardiac glycosides have narrow therapeutic indices and their use is frequently accompanied by toxic effects that can be severe or lethal. The most important toxic effects, in terms of risk to the patient, are those that involve the heart (e.g., abnormalities of cardiac
30 rhythm and disturbances of atrio-ventricular conduction). Gastrointestinal disorders, neurological effects, anorexia,

-3-

blurred vision, nausea and vomiting are other common cardiac glycoside-induced reactions.

In the case of ischemic heart disease, other classes of compounds have been used to improve blood flow to the failing myocardium through dilation of the coronary arteries. Examples of these agents include nitrates, such as isosorbide dinitrate, and calcium channel blockers such as diltiazem, nifedipine and verapamil.

Nitrates and calcium channel blockers do not have direct effects on the contractile mechanism of the cardiac muscle cells. That is, they are not positive inotropic agents. Digitalis and ouabain, on the other hand, are strong inotropes, but do not have vasoactive (dilatatory) effects on the coronary circulation. Hypothalamic inhibitory factor (HIF) has also been shown to produce a positive inotropic effect on cardiac muscle cells (U.S. patent application No. 08/338,264, filed November 10, 1994). However, HIF was not shown to have vasoactive (dilatatory) effects on the coronary circulation.

A single compound which has both inotropic effects and coronary vasodilatory effects would be highly desirable as an agent to treat ischemic heart failure since both abnormalities are present and each contributes to the morbidity of the condition. To this point, such a compound has not been identified or made available pharmaceutically for the treatment of ischemic heart failure.

SUMMARY OF THE INVENTION

The present invention relates to a method for increasing coronary perfusion in a mammalian host comprising administering to the host in need thereof an

-4-

effective amount of hypothalamic inhibitory factor (HIF).
The invention also relates to a method for producing an
increased coronary vasodilatory effect in a mammalian host
comprising administering to a mammalian host in need
5 thereof an effective amount of HIF.

In a particular embodiment, the invention pertains to
a method for preventing or treating ischemic cardiac
malfunction comprising administering to a mammalian host in
need thereof an effective amount of HIF. The invention
10 also relates to a method for preventing or treating
coronary artery restenosis comprising administering to a
mammalian host in need thereof an effective amount of HIF.
Also encompassed by the present invention is a method for
treating stenosis of a coronary artery in a mammalian host
15 comprising administering to the host in need thereof an
effective amount of HIF. Stenosis of a coronary artery can
be due to, for example, atherosclerosis and/or coronary
artery restenosis following the intervention of mechanical
dilatation or certain bypass graft operations.

20 HIF slows heart rate in spontaneously beating cardiac
myocytes, further reducing cardiac work and thereby
diminishing oxygen requirement. The invention provides a
single compound, HIF, which has the combined properties of
producing a positive inotropic effect and enhancing
25 coronary flow, and which can be used prophylactically
and/or therapeutically to treat the identified conditions
associated with heart failure.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the unexpected
30 discoveries that hypothalamic inhibitory factor (HIF) has a

-5-

positive inotropic effect in a whole organ preparation, such as an isolated perfused heart, and that HIF, to a greater extent than ouabain, dilates the coronary arteries in the isolated perfused heart as manifested in increased coronary flow in hearts treated with HIF. As defined herein "coronary flow" and/ or "coronary perfusion" refers to milliliters per minute of physiologic buffer perfusing the coronary arteries. As demonstrated herein HIF causes both enhanced contractility and simultaneous increase in coronary perfusion when administered to an intact heart. The present invention relates to methods for increasing coronary perfusion in a mammalian host comprising administering to the host in need thereof an effective amount of HIF. The present invention also relates to a method of producing a positive inotropic effect and an increased vasodilatory effect in a mammalian host comprising administering to the host in need thereof an effective amount of HIF. The mammalian host can be any mammal which is in need of increased coronary perfusion, and includes, for example, human, canine, feline, bovine and murine hosts.

HIF is an endogenous inhibitor of $\text{Na}^+\text{-K}^+\text{-ATPase}$, which has been isolated from bovine hypothalamus and human plasma, and structurally characterized as an isomer of the plant cardiac glycoside, ouabain (Tymiak, A.A., et al., *Proc. Natl. Acad. Sci., USA*, 90:8189-8193 (1993); Zhao, N., et al., *Biochemistry*, 34:9893-9896 (1995)). HIF for use in the present invention can be obtained by purifying HIF from natural sources or chemically synthesizing HIF. In a preferred embodiment, purified HIF is used. Purified HIF refers to HIF which is substantially free of or isolated

-6-

from other tissue or fluid protein components and contaminants. Various procedures may be used to purify HIF from natural sources. For example, as described in U.S. patent application No. 08/338,264, filed November 10, 1994, which is incorporated by reference, HIF has been purified to homogeneity using an affinity chromatography method in which purified renal Na⁺, K⁺-ATPase (e.g. isolated from a canine) is coupled to paramagnetic particles through a glutaraldehyde bridge (Tymiak, A.A., et al. *Proc. Natl. Acad. Sci., USA*, 90:8189-8193 (1993)). The enzyme immobilizes bound HIF in the presence of Mg⁺⁺ and inorganic phosphorous with high affinity, and after washing away contaminating materials, HIF is eluted from the affinity column by chelating Mg⁺⁺ with EDTA. A subsequent HPLC step results in purification of HIF (Tymiak, A.A., et al. *Proc. Natl. Acad. Sci., USA*, 90:8189-8193 (1993)). HIF has also been purified to homogeneity using an immunoaffinity chromatography method in which an antibody which binds to HIF is coupled to the resin of an immunoaffinity column. The antibody recognizing HIF immobilizes bound HIF with high affinity, and after washing away contaminating materials, HIF is eluted from the affinity column. A subsequent HPLC step can be used to further purify HIF. See U.S. patent application No. _____, filed May 30, 1997, Attorney Docket No. BION97-01, which is incorporated by reference. It should be noted that pharmaceutically acceptable salts of HIF are also contemplated. Suitable salts include those well known to those of skill in the art.

As defined herein, an "effective amount" is an amount sufficient, when administered to the host, to result in an increased coronary vasodilatory effect and, preferably, a

-7-

positive inotropic effect also in the mammalian host relative to the inotropic and coronary vasodilatory effects when an effective amount of HIF is not administered. These effects are associated with increased coronary perfusion and can be utilized to treat and/or prevent ischemic cardiac malfunction, coronary artery restenosis and stenosis of a coronary artery in a mammalian host. The amount of HIF used to treat a host will vary depending on a variety of factors, including the size, age, body weight, general health, sex and diet of the host. In addition, an effective amount of HIF will depend on the nature of the disease, and can be determined by standard clinical techniques. The precise dose to be employed will also depend on the route of administration and the seriousness of the cardiac disease or disorder, and should be decided according to the judgement of the practitioner and each patient's circumstances. However, suitable dosage ranges for intravenous administration are generally about 4.5 ug to about 20 ug per kilogram body weight. In one embodiment the dosage is about 4.5 ug per kilogram body weight, in another embodiment the dosage is about 12 ug per kilogram body weight, and in a further embodiment the dosage is about 20 ug per kilogram body weight. Suitable dose ranges for oral administration of HIF are the same as for intravenous administration of HIF, but the HIF dosage is administered orally over several days. Effective doses may be extrapolated from dose response curves derived from *in vitro* or animal test models.

The HIF of the present invention can be administered prophylactically to a host as a method of preventing the conditions described herein. Alternatively, HIF can be

-8-

administered therapeutically to a host as a method of treating an existing disease and/or condition in the host, and can result in amelioration or elimination of the disease and/or condition.

5 The formulation and route of delivery of HIF to the host can be accomplished in a variety of ways. Routes of administration include intradermal, transdermal (e.g. slow release polymers), intramuscular, intraperitoneal, intravenous, intracardiac, subcutaneous, oral, epidural, 10 and intranasal routes. Any other convenient route of administration can be used, for example, infusion or bolus injection, or absorption through epithelial or mucocutaneous linings. The HIF can be administered together with other components or biologically active 15 agents, such as pharmaceutically acceptable surfactants (e.g., glycerides), carriers (e.g., saline, buffered saline, dextrose, water, glycerol, ethanol), excipients (e.g., lactose), diluents and vehicles and combinations thereof. The formulation should suit the mode of 20 administration.

The composition can also include, if desired, minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained 25 release formulation or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium 30 saccharine, cellulose, magnesium carbonate, etc.

-9-

In a preferred embodiment, the HIF is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to the mammalian host (e.g., a human). For example, HIF for
5 intravenous administration can be a solution in sterile isotonic aqueous buffer. Where necessary, the HIF composition may also include a solubilizing agent and a local anesthetic to ease pain at the site of injection. Generally, the ingredients are supplied either separately
10 or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampule or sachette indicating the quantity of active agent. Where the composition is administered by injection, an ampule of
15 sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration. Thus, the invention also relates to the use of HIF in the manufacture of medicaments for the treatment and/or prevention of ischemic cardiac malfunction, coronary artery
20 restenosis and/or stenosis of a coronary artery.

The invention can be used to prevent and/or treat a patient with ischemic heart failure, which is due, for example, to stenosis of the coronary artery. It is estimated that four million Americans suffer from heart
25 failure, and that 60-70% of these cases (2.4-2.8 million) are on the basis of ischemic disease. The combined advantages over digitalis and ouabain of less toxicity and enhanced coronary perfusion makes the invention a therapeutically desirable substitute for digitalis in these
30 patients.

-10-

The invention also relates to methods wherein HIF is administered to prevent and/or treat coronary artery restenosis in patients who have undergone coronary artery angioplasty or certain kinds of coronary artery bypass surgery for conditions such as ischemic heart disease.

HIF does not manifest the same toxicity profile as the cardiac glycosides. Therefore, prevention and/or treatment of conditions associated with heart failure, such as ischemic cardiac failure and coronary artery restenosis, with HIF can be accomplished with less risk of toxicity to the patient, and both abnormal muscular contractility and inadequate flow of blood and oxygen to the ischemic myocardium can be concurrently treated and/or prevented with a single compound.

This invention is illustrated further by the following exemplification, which is not to be construed as limiting in any way.

EXEMPLIFICATION

Measurement of positive inotropic effects and coronary flow

Guinea pigs were anesthetized with 65-100 mg pentobarbital in the peritoneal cavity followed by anticoagulation using 200 units of heparin via femoral vein. Sternotomy was performed, the beating heart removed and placed in a bowl containing room temperature saline. The aorta was trimmed and cannulated onto the perfusion apparatus. Retrograde perfusion of Kreb's solution (37°C) at 70 mm Hg bubbled with 95% O₂/5% CO₂ from a 3 liter reservoir was begun within one minute of cardiectomy. A micro flexible temperature probe was inserted in the open

-11-

pulmonary artery allowing coronary effluent to drain. The left atrial appendage was incised, the left ventricle vented, and a water-filled balloon-tipped catheter inserted into the left ventricle chamber. Left ventricular systolic pressure (LVP), left ventricular end diastolic pressure (LVEDP), maximal rate of rise of left ventricular pressure (dp/dt), and aortic root pressure (AOP) were recorded on a precalibrated multichannel dynograph. Coronary flow (CF) is measured by timed volumetric collection. The heart was paced (HR) at 272 BPM by way of electrodes attached to the right atrium. After 30 minutes of perfusion, coronary flow was then diverted into a recirculating perfusion circuit primed with 60 cc of Kreb's solution.

The circuit was started with the coronary effluent draining into a reservoir. The effluent was pumped out the reservoir through a 5 micron filter that passed through a low volume silastic oxygenator and up a pressure column. The oxygenated effluent flowed back to the aorta or out an overflow column that drained to the reservoir. Pump output was kept higher than the coronary flow allowing a constant aortic root pressure. Heart temperature (HT), pH, O_2 and CO_2 of the perfusate were held constant by water jacketing the oxygenator and perfusion circuit and no additional Kreb's solution needed to be added during the recirculation intervals. During the first ten minutes of recirculating, the intraventricular balloon volume was adjusted to reach an LVEDP of 10 mm Hg. The heart was switched back to non-recirculation for ten minutes while the recirculating circuit was emptied and refilled with fresh Kreb's solution. Two additional recirculation non-recirculation cycles were carried out with LVEDP adjusted and parameters

-12-

measured. LVP, LVEDP, dP/dT, AOP, HR, HT and CF were recorded at 5 minute intervals for 15 minutes. Then a coded aliquot of ouabain, HIF or placebo was added to the recirculating Kreb's solution. Parameters continued to be
5 measured at 5 minute intervals for 30 additional minutes.

Results

HIF has been previously shown to produce positive inotropic effects in isolated, spontaneously beating myocytes in culture, with greater potency and a different
10 toxic profile from ouabain (Hallaq, H. and Hauptert, G., Jr., *Proc. Natl. Acad. Sci., USA*, 86:10080-10084 (1989)). As shown herein, HIF also produces a positive inotropic effect in an intact, whole organ preparation. 20 µg HIF was prepared to homogeneity as confirmed by mass spectral
15 measurement, for infusion into an isolated, perfused guinea pig heart preparation according to the Langendorf methodology (Hendren, W.G., et al., *J. Thor. Cardiovasc. Surg.*, 94:614-625 (1987)). This methodology allows continuous measurement of left ventricular pressure
20 generation, the first derivative of the pressure measurement (dP/dT), end diastolic pressure and coronary flow. Pilot studies with ouabain were carried out to determine the minimal consistently effective dose to produce an enhanced inotropic effect. For both ouabain and
25 HIF, this amount was 4 µg of pure inhibitor administered to a perfusion volume of 60 ml Krebs-Heinsleit buffer (final concentration 1×10^{-7} M). Coded samples were prepared of HIF, ouabain and placebo (vehicle), and administered randomly to the preparation. The one conducting the
30 experiments was blinded to the samples.

-13-

Results are shown in the Table. The Table shows the effects of HIF, ouabain and placebo on performance characteristics and coronary flow in isolated, perfused guinea pig heart. Both HIF and ouabain showed significant increases in left ventricular pressure (lvp) and rate of development of pressure (dP/dT) compared to the preadministration control period, indicating positive inotropic response. Placebo produced no change. HIF and ouabain also produced a statistically significant increase in coronary flow, but the percent increase from baseline for HIF surprisingly was two-fold greater than the increase produced by ouabain (Table). Thus, the mammalian-derived HIF was a significantly more potent coronary vasodilator than the plant-derived ouabain. HIF produced a statistically significant increase in coronary flow and is less toxic than ouabain.

-14-

TABLE 1: Effects of HIF, ouabain and placebo on performance and perfusion characteristics in isolated, perfused guinea pig heart (LVP, left ventricular pressure; dP/dt, rate of development of pressure)

GROUPS	LVP	LVP post intervention	%	dPdt mmHg.sec-1	dP/dt post intervention	%	End Dia pressure	EDP post intervention	%	coronary flow cc/min	CF post intervention	%
ouabain n=9	73.82	90.44 ^{^*}	22.5	2326	2856 ^{^*}	22.7	7.66	8.01	4.6	25.84	27.77 [*]	7.5
placebo n=8	67.17	68.29	1.7	2104	2088	-0.7	7.77	9.19	18.2	25.89	25.09	-3.1
HIF n=9	70.54	89.91 ^{^*}	27.4	2292	2884 ^{^*}	25.8	7.6	7.77	2.1	24.37	27.82 [*]	14.1

[^] P<0.01 intervention vs placebo

^{*} P<0.01 pre intervention vs post intervention

-15-

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific embodiments of the
5 invention described herein. Such equivalents are intended to be encompassed in the scope of the following claims.

-16-

CLAIMS

We claim:

1. A method for increasing coronary perfusion in a mammalian host comprising administering to the host in need thereof an effective amount of hypothalamic inhibitory factor.
2. The method of Claim 1 wherein the mammalian host is a human.
3. The method of Claim 1 wherein the hypothalamic inhibitory factor is administered intravenously.
4. A method for producing an increased coronary vasodilatory effect in a mammalian host comprising administering to a mammalian host in need thereof an effective amount of hypothalamic inhibitory factor.
5. The method of Claim 4 wherein the mammalian host is a human.
6. The method of Claim 4 wherein the hypothalamic inhibitory factor is administered intravenously.
7. A method for preventing or treating ischemic cardiac malfunction comprising administering to a mammalian host in need thereof an effective amount of hypothalamic inhibitory factor.

-17-

8. The method of Claim 7 wherein the mammalian host is a human.
9. The method of Claim 7 wherein the ischemic cardiac malfunction is due to stenosis of the coronary artery.
- 5 10. The method of Claim 7 wherein the hypothalamic inhibitory factor is administered intravenously.
11. A method for preventing or treating coronary artery restenosis comprising administering to a mammalian host in need thereof an effective amount of
10 hypothalamic inhibitory factor.
12. The method of Claim 11 wherein the mammalian host is a human.
13. The method of Claim 11 wherein the hypothalamic inhibitory factor is administered intravenously.
- 15 14. A method for treating stenosis of a coronary artery in a mammalian host comprising administering to the host in need thereof an effective amount of hypothalamic inhibitory factor.
15. The method of Claim 14 wherein stenosis of the
20 coronary artery is due to coronary artery restenosis.
16. The method of Claim 14 wherein the mammalian host is a human.

-18-

17. The method of Claim 14 wherein the hypothalamic inhibitory factor is administered intravenously.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/10890

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/705

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 433 074 A (GEN HOSPITAL CORP) 19 June 1991 cited in the application see abstract; claims 1-3 ---	7-10
A	N. ZHAO ET AL.: "NA,K-ATPase INHIBITORS FROM BOVINE HYPOTHALAMUS AND HUMAN PLASMA ARE DIFFERENT FROM OUABAIN: NANOGRAM SCALE CD STRUCTURAL ANALYSIS" BIOCHEMISTRY, vol. 34, no. 31, 1995, pages 9893-9896, XP002081420 cited in the application see the whole document --- -/--	1-17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 October 1998

Date of mailing of the international search report

02/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 98/10890

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TYMIAK A A ET AL.: "PHYSICOCHEMICAL CHARACTERIZATION OF A OUABAIN ISOMER ISOLATED FROM BOVINE HYPOTHALAMUS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 90, no. 17, 1 September 1993, pages 8189-8193, XP002074993 cited in the application see the whole document</p> <p>---</p>	1-17
A	<p>R. KOHN ET AL.: "ENDOGENOUS DIGITALIS-LIKE FACTOR IN PATIENTS WITH ACUTE MYOCARDIAL INFARCTION" COR ET VASA, vol. 34, no. 3, 1992, pages 227-237, XP002081545 see the whole document</p> <p>---</p>	1-17
A	<p>P. DELVA ET AL.: "INCREASE IN PLASMA DIGITALIS-LIKE ACTIVITY DURING PERCUTANEOUS TRANSLUMINAL CORONARY ANGIOPLASTY IN PATIENTS WITH CORONARY STENOSIS" LIFE SCIENCE, vol. 47, no. 5, 1990, pages 385-389, XP002081546 see the whole document</p> <p>-----</p>	1-17

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 10890

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 1-17
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/10890

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0433074 A	19-06-1991	AT 114475 T	15-12-1994
		CA 2032079 A	14-06-1991
		DE 69014569 D	12-01-1995
		DE 69014569 T	13-07-1995
		DK 433074 T	08-05-1995
		ES 2069030 T	01-05-1995
		GR 3015201 T	31-05-1995
		JP 4054128 A	21-02-1992
		US 5716937 A	10-02-1998
